<u>Photo</u>	Botanical Name	Comm on Name	Annual or Perenn ial	Heigh t	Sprea d	Rate of Growth	Climat e	Water/ Soil	Hardine ss Zone	<u>Info</u>	<u>Foliage</u>	Flowers/ Fruits Bloom/ Seed Times	Eco- Sculpture	Vend or	Eco thoughts	Medicinal Properties	Nutritional (edible/ culinary Nutritional value/ edible parts)	Pests or Disease S	Poisonous Parts/ indication s	Wildlife Value
	Ajuga reptans (x tenorii) 'Chocolate Chip'	Buglew	Perenni	10-15 cm 4-6 inches	30-45 cm 12-18 inch	Medium Not as quick to spread as many of the larger selectio ns	shady area	reasonabl y moist soil	Zone 3		unusually narrow foliage is dark green, overlaid with chocolate- brown	Short spikes of showy blue flowers appear in late spring	Bruce's Eagles, Salmon, Pandas 2009	Nat's	Good coverage, nice dark color. Needs flower heads removed regularly.	Has a long history of use as a wound herb and, although little used today, it is still considered very useful in arresting haemorrhages and is also used in the treatment of coughs and spitting of blood in incipient consumption. The plant contains digitalis-like substances (these are commonly found in Digitalis species and are used in treating heart complaints) and is thought to possess heart tonic properties. It has also been considered good for the treatment of excessive alcohol intake. The whole plant is aromatic, astringent and bitter. The plant is usually applied externally. It is harvested as it comes into flower in late spring and dried for later use. It is also commonly used fresh in ointments and medicated oils. A homeopathic remedy is made from the whole plant. It is widely used in various preparations against throat irritations and especially in the treatment of mouth ulcers.	salads.	susceptib le to crown rot.	The plant is said to be a narcotic hallucinoge n that is known to have caused fatalities	Attracts butterflies and hummingbi rds, the flowers being very attractive to bees

Alternant era dentata 'Purple Knight'	Joseph' s coat, Joy weed Calico plant	Annual	6-12 in 15-30 cm (12-18 in 30-45 cm)		fast Full Sun to Partial Shade	Water: Medium Soil type: Well- drained light	Perenni al in Zone 10 to 11	The. Give "Purple Knight" a lot of space - it grows big and fast, up to 3' tall and 3' across.	Grown for purple stems that spread out wide, loaded with glistening, two-toned foliage - purple on top, burgundy below		Horses 2009	Mead owlan ds	unknown	unknown	Spider mites and Fusarium wilt may be problems	unknown	This plant is attractive to bees, butterflies and/or birds
Alternant era 'True yellow'	Calico Plant	Annual	6-12 in 15-30 cm	4-10 in 10-15 cm	Full Sun to Shade Prefer s part sun	Average - Moist well drained soil	Perenni al in Zone 9-11		Solid yellow green foliage holds up great in the heat			Mead owlan ds			Mealybu gs and Rhizocto nia can be a problem if the plant is stressed. Aphids and white flies can be controlle d with good airflow and drainage.		
Alternant era ficoidea 'Red carpet'	Joseph' s Coat	Annual	12-14 inches	15-18 in. (38-45 cm) 18-24 in. (45-60 cm)	Full sun to part shade	Average water needs; do not overwater	Perenni al in zones 8-10.		This one is grown for its colourful foliage. Beautiful leaves with reds, yellows, burgundy and greens	Inconspicu ous white		Mead owlan ds	unknown		Spider mites and Fusarium wilt may be problems	Danger: N/A	

	Antennaria dioica 'Rubra'	Pink Pussy toes	Perenni	10-15 cm 4-6 inches	cm 10-12		especi ally suited to hot sunny locatio ns	A tough plant will take poor, dry soil. Very drought tolerant.	Zone 5	a very low, creeping mat of tiny, bright silvery-grey leaves.	Short stems of fuzzy pink flowers appear from June to July. The plant is self-fertile.	Bruce's Eagles Pandas 2009	Nice low mat forming	Has been little used in herbal medicine though it was once used in mixtures for the treatment of bronchitis and bilious conditions. The whole plant is antitussive, astringent, cholagogue, discutient, diuretic and emollient. The plant is very rich in mucilage which makes it very valuable in the treatment of chest complaints. It is also used in the treatment of liver and gall bladder complaints, hepatitis and diarrhoea. Externally it is used as a gargle for treating tonsillitis and as a douche for vaginitis. The herb is gathered in May before it comes into flower and can be dried for later use.		Root, stem, crown and collar rots		tracts utterflies
--	---------------------------------	-----------------------	---------	------------------------------	-------------	--	--	--	--------	---	--	-------------------------------------	----------------------	---	--	-----------------------------------	--	----------------------

n p	Ophiopogo n olaniscapu s nigrescens	Black Mondo Grass	evergre en perenni al	Its strap-like leaves will reach 15 inches (37.5 cm) tall	Leave s will be up to ¼ inch (6 mm) wide.		ant of full sun.	Moist well-drained soil. In container s in the greenhou se, we use a soil mix consistin g of 2 parts peat moss to 2 parts loam to 1 part sand or perlite will with a During the growing season the plants are kept moist for optimum growth and fertilized every other week with a balanced fertilizer diluted to ½ the strength recomme nded on the label	Zones 5-9. Only winter hardy to USDA Zone 7 (roots will usually be killed when tempera tures dip below 10 F),	a great groundc over or edge plant for the border of a garden. and Plants	The foliage starts out green and then quickly turns dark black.	Blooming Time: In late summer to early fall, the plants bloom, forming white flowers with a pink flush 1/4 inch (0.6 cm) flowers. Shiny black fruits follow flowers.	Crane		No serious insect or disease problems	

Arctostaph ylos uva- ursi	Bearber ry or Kinnikin nick	Perenni	10-15 cm 4-6 in	30-60 cm 12-23 inch	to	ade no control ade state	Best in a well-drained sandy to loam soil, this dislikes poorly-drained clay. Drought tolerant, once establish ed.	Zone 4	This is a selection of a native North American wildflower, introduced by the U.B.C. Botanical Gardens. Will tolerate light foot traffic.	particularly in	bell- shaped flowers appear in spring, followed by	Salmon	Nat's	Doesn't bulk up enough, stems tend to be too long. Most of it died off.	range of complaints and has also been used in conventional herbal medicine for hundreds of years, it is one of the best natural urinary antiseptics. A tea	Fruit can be added to stews etc, it is a good source of carbohydra tes, can also be used to make a cooling drink or used for preserves etc. A tea is made from the dried leaves.	
Armeria maritima	Pink Thrift	Perenni al	10-15 cm 4-6 inches	15-30 cm 6-12 inch	loo n. ca gr in	catio t	Can tolerate drought.	Zone 2		Plants form a low tufted mound of grassy green leaves.	This selection bears deep pink pompon flowers for many weeks starting in late spring, and usually repeat flowers in the fall.	Dragster scoop		The flowers looked good, too dry a location for it, though and was removed.	The dried flowering plant is antibiotic and has been used in the treatment of obesity, some nervous disorders and urinary infections.	Leaves and roots are edible	Pollinated by Bees, flies, beetles, Lepidopter a (Moths & Butterflies). It is noted for attracting wildlife.

Cornus Canadensi s	Bunchb erry, Creepin g dogwoo d	Perenni	10-15 cm 4-6 inches	cm 8-12	Slow	moist shady area.	Prefers a damp acidic soil. It can grow in semishade (light woodland) or no shade. It requires moist soil. Grows well in heavy clay soils. Grows best in sandy soils.	Zone 2	This woodla nd plant is one of our most attractive vildflowers.	colour. Plants go completely dormant in	Single white Dogwood flowers appear in late spring, followed by a colourful crop of scarlet berries in the fall.	Salmon, Crane egg	Most of it died off.	The leaves and stems are analgesic, cathartic and febrifuge. A tea has been used in the treatment of aches and pains, kidney and lung ailments, coughs, fevers etc. A strong decoction has been used as an eye wash. The fruits are rich in pectin which is a capillary tonic, antioedemic, anti-inflammatory, antispasmodic and hypotensive. Pectin also inhibits carcinogenesis and protects against radiation. A tea made from the roots has been used to treat infant colic. The mashed roots have been strained through a clean cloth and the liquid used as an eyewash for sore eyes and to remove foreign objects from the eyes.	Fruit - raw or cooked		
Festuca rubra	Red fescue	Perenni	10-80 cm 24 inches		Medium	Prefer s sun to part shade.	Has good drought and excellent cold tolerance.	Zone 3	Native to the U.S. It is used as a cool season turf grass and groundc over.	This is fine-leaved turf grass.	Non flowerin g	Crane Nat's	Really poor coverage over summer and winter. Dormant/ dead looking and not sky reaching. Thatch got into all soil, and we had to remove and redo all stuffing etc on crane.		Seeds are edible. No further details but the seed is rather small and would be very fiddly to utilise.	It is very susceptib le to summer diseases in hot climates.	It is a nutritious and palatable forage grass for domestic and wild animals.

Feet nut case	Gaultheria shallon	Salal	Perenni	60-12 0 cm 23-47 inch	45-90 cm 18-35 inch	Medium	Partial shade to full shade preferr ed	Requires acidic soil. Drought tolerant once establish ed	Zone 5	An evergre en shrub native to the Pacific Northw est. A purple dye is obtaine d from the fruit. A greenis hyellow dye is obtaine d from the infused leaves.	thick patch of large green, leathery leaves, and these are still used as commercial florist's greens. This compact,		Salmon, Crane egg	Nat's	Much too big for our sculptures we wanted G. procumben s (wintergree n) None surviving.	A poultice of the toasted, pulverized leaves has been applied to cuts. A poultice of the chewed leaves has been applied to burns and sores. The leaves have been chewed to dry the mouth. An infusion of the leaves have been used as a stomach tonic and a treatment for diarrhoea, coughs, TB etc.	Fruit - raw, cooked or dried for later use, it is sweet and juicy with a pleasant flavour it makes good raw eating. The fruit can also be made into preserves, pies, drinks etc or be dried and used like raisins. A pleasant tea is made from the leaves.	Black mildew, powdery mildew, leaf gall, fungal spots.	
	Iberis sempervire ns	Evergre en candytu ft	Perenni	20-25 cm 8-10 inch	30-90 cm 12-35 inch	Medium	Droug ht toleran t, once establi shed.	Prefers a well-drained site, so avoid heavy clay soils that stay wet in winter.	Zone 3	Not easily divided.	Numerous, oblong, entire, dark green, leathery leaves (1-1.5" long).	Spring blooming, with loads of goodsized white flowers for several weeks.	Crane egg		Does better in sculpture than in trays			Crown rot can devastat e plantings. Suscepti ble to club root which results in stunted growth. Desiccati on and sun scorch may damage the evergree n foliage in cold winter climates.	

	Isotoma fluviatilis White	White Star Creeper	Perenni	2-4 cm 1-2 inches	15-30 cm 6-12 inch	Medium		Average to moist well- drained soil	Zone 6	Plants are easily ripped apart into small pieces to make new planting s, and this should be done in early spring. Evergre en.	flat carpet of tiny green leaves,	smothered with starry, white flowers all summer long	Bruce's Eagles	Really good coverage, nice white flowers. But is invasive and will probably take over whole sculpture.	No seriou insect diseas proble . Water for aphids	or may cause skin and/or seye h irritation in some	
-SHEPNIS NOOD	Leptinella squalida 'Platt's Black'	Black Brass Buttons	Perenni	2-5 cm 1-2 inches	25-30 cm 10-12 inch	Medium	part shade. Tolerat es light shade, particu larly in hot summ er	Easy to grow in any soil that does not completel y dry out. Well-drained soils in. Intoleran t of drought.	Zone 4	fall.	(to 2" long and to 1/2" wide) are obovate. Foliage will tolerate some foot	It features button-like yellow -Dark gold flowers bloom in early summer, and are followed by tiny fruit. Neither the flowers nor the fruit are considered to be particularly ornamental .	Bruce's Eagles	Looks crispy and dried out, but is surprisingly still alive. Nice dark color for contrast.	No seriou insect diseas proble. Som susce ility to vertici m wilt leaf sy and si rots. Aphid mites and le miner are occas al visitor	or see ms se potib liu soots eem se, af se soon	

Santolina chamaecyp arissus	Cotton Lavend er	Perenni	0.4 - 0.7m	0.3 - 0.6m	Fast	wants full sun	Very drought tolerant	Zone 6	A branchi ng aromati c (camph or-like fragran ce) Mediter ranean shrub.	Woolly stems and soft feathery silver-grey leaves. Leaves are 1mm, arranged geometrically	Flower is a 1cm head of yellow disk florets in summer. Flowers have an unpleasant smell.			The leaves and flowering tops are antispasmodic, disinfectant, emmenagogue, stimulant and vermifuge. Cotton lavender is rarely used medicinally though it is sometimes used internally as a vermifuge for children and to treat poor digestion and menstrual problems. When finely ground and applied to insect stings or bites, the plant will immediately ease the pain. Applied to surface wounds, it will hasten the healing process by encouraging the formation of scar tissue. The leaves and flowering stems are harvested in the summer and dried for later use.	The aromatic leaves are used as a flavouring for broths, sauces, grain dishes.	Soggy soil in winter can cause root rot.	The bruised leaves have been known to cause a severe rash on sensitive skins	
Sedum acre	Gold moss Stonecr op	Perenni	5-8 cm 2-3 inches	30-60 cm 1-2 feet	Vigorous	not thrive in dense shade with limited	dry sandy soil, grows well in poor soils, sand, rock gardens, and rich dirt	Zone 2	Native to Europe, but also naturali sed in North Americ a. Evergre en.		Starry yellow clusters of flowers appear in June-July.	Salmon	looks nice	The herb is astringent, hypotensive, laxative, rubefacient, vermifuge and vulnerary. It is considered to be a useful medicinal plant by some herbalists, though others do not use it because of the violence of its operation when taken internally. One of its best uses is as an effective and harmless cornremover, it can also be used to bring boils to a head, though this can also cause some local irritation. The bruised fresh plant is applied as a poultice to wounds and minor burns.	Leaves - raw or cooked. Rich in vitamin C, but it has a bitter acrid taste.	No serious insect or disease problems	The sap can cause blisters or irritate the skin of some people. Large quantities can cause stomach upsets. It is best to dry the leaves (which can be difficult because they are very fleshy) and then powder them and use them to add a peppery taste to foods	pollinated by Bees, flies.

Sedum hybridum 'Immergrun chen'	Stonecr		15-30 cm 6-12 inch	30-38 cm 12-15 inch			Drought tolerant		Nice fall and winter color. Evergre en.	foliage on a compact low growing	Flowers are pale yellow.	Salmon	Good green coloring, nice matting. Tends to fall, as new growth is longer.		
Sedum kamtschati cum	Stonecr	Perenni	6 inches	10 inches		Will handle some shade.	drought tolerant	Zone 4	This is not a NA native. Decidu ous; trim stems and dead flower heads back in early spring.	A somewhat taller and fleshy Sedum	summer blooming, star- shaped yellow flowers open from pink buds and mature to crimson	Panda	Deciduous! !!! Won't work through winter. Removed from Panda!	Mealybu gs, scale, slugs, snails	
Stipa tenuissima	Mexica n Feather Grass	Someti mes acts as a self- seeding annual	45-60 cm 18-23 inch	30-45 cm 12-18 inch	Fast	sunny	Drought tolerant once establish ed		A clump- forming grass, this is a terrific accent for the front of a border, the rock garden, or in mixed contain ers.	Plants form a mound of fine, hair-like green leaves.	bearing pale yellow flowers that turn into whiskered tan- coloured heads by late summer. Very soft and silky in appearanc e.	Parsley and Sage	Great Mane material for horses! Self seeds though, and really fast growing.		